



Extension


UNIVERSITY OF WISCONSIN-MADISON

Provided to you by:

Impatiens Necrotic Spot

Ann Joy and Brian Hudelson, UW-Madison Plant Pathology

What is impatiens necrotic spot? Impatiens necrotic spot is a viral disease that causes considerable losses to greenhouse-grown ornamentals and, to a lesser extent, vegetable crops. In the southern United States, impatiens necrotic spot can also be a problem on field crops.



Ringspots on coleus leaf caused by impatiens necrotic spot virus. (Photo courtesy of Margaret Daughtrey)

Ornamental crops affected by impatiens necrotic spot include impatiens, gloxinia, cineraria, cyclamen, exacum, petunia, begonia, primrose and ranunculus. Susceptible vegetables include cucumber, tomato and pepper. Impatiens necrotic spot also affects weeds including chickweed, jewelweed, oxalis, and gill-over-the-ground.

What does impatiens necrotic spot look like?

Symptoms vary from species to species and from cultivar to cultivar. Environmental conditions can also influence symptom development. Infected plants may exhibit chlorotic or necrotic spotting; stem, vein or growing point necrosis; ringspots, mosaic or line patterns on leaves; color break in

flowers; wilting and collapse. Plants that are infected at a young age tend to exhibit more severe symptoms than those infected at an older age. Some plants may remain symptomless but still can be a source of the causal virus, leading to infection other plants.

Where does impatiens necrotic spot come from? Impatiens necrotic spot is caused by the *Impatiens necrotic spot virus* (INSV). The primary means of plant-to-plant spread of INSV is by the western flower thrips (*Frankliniella occidentalis*). This insect acquires INSV when feeding on infected plant sap, carries the virus to its next feeding site, and transmits the virus to the new plant after 15 to 30 minutes of feeding. Only a few thrips are needed to spread INSV rapidly throughout a greenhouse. Numbers of INSV-infected plants also can be increased simply by taking vegetative cuttings from infected plants.

How do I save a plant with impatiens necrotic spot? After infection, INSV spreads throughout a plant. Therefore plants remain infected indefinitely. Infected plants cannot be treated to remove the virus and should be destroyed by burning (where allowed by local ordinance), burying or composting.



Extension

UNIVERSITY OF WISCONSIN-MADISON

How do I control impatiens necrotic spot in the future? Impatiens necrotic spot control focuses on excluding INSV and preventing its spread. Inspect any plants entering a greenhouse (e.g., new plant shipments, plants moved in from outdoors) for viral symptoms and thrips. Isolate new plants until they are



Symptoms of impatiens necrotic spot on Gloxinia. (Photo courtesy of Margaret Daughtrey)

determined to be thrips- and virus-free. In commercial settings, isolate breeding and stock plants from production, and do not carry over plants from fall production into the spring. Destroy any symptomatic plants. Control greenhouse weeds, which can be symptomless reservoirs of INSV. Monitor and control thrips populations. Exclude thrips by screening greenhouse vents and doors (screening should have apertures that are ≤ 0.135 mm). Set up yellow, white or blue sticky traps to monitor for thrips in growing areas and near greenhouse entry points. Indicator plants such as *Petunia x hybrida* 'Summer Madness', 'Super Magic Coral', or 'Red Cloud' (with flowers removed) can provide early warning signs of a problem with thrips.

Petunia leaves on which INSV-free thrips feed will develop whitish feeding scars, while those leaves on which INSV-carrying thrips feed will develop small brown to black spots, turning tan with a dark border. See also University of Wisconsin Garden Facts XHT1022, *Managing Thrips in Greenhouses*, for additional pointers on thrips control, including insecticide recommendations.

For more information on impatiens necrotic spot: Contact the University of Wisconsin Plant Disease Diagnostics Clinic (PDDC) at (608) 262-2863 or pddc@wisc.edu.

© 2005-2024 by the Board of Regents of the University of Wisconsin System doing business as the Division of Extension of the University of Wisconsin-Madison.

An EEO/Affirmative Action employer, University of Wisconsin-Madison provides equal opportunities in employment and programming, including Title IX and ADA requirements. This document can be provided in an alternative format by calling Brian Hudelson at (608) 262-2863 (711 for Wisconsin Relay).

References to pesticide products in this publication are for your convenience and are not an endorsement or criticism of one product over similar products. You are responsible for using pesticides according to the manufacturer's current label directions. Follow directions exactly to protect the environment and people from pesticide exposure. Failure to do so violates the law.

Thanks to Tom German, Andy Witherell and Anna Whitfield for reviewing this document.

A complete inventory of University of Wisconsin Garden Facts is available at the University of Wisconsin-Madison Plant Disease Diagnostics Clinic website: <https://pddc.wisc.edu>.

Submit additional lawn, landscape, and gardening questions at <https://hort.extension.wisc.edu/ask-a-gardening-question/>.