



Extension

UNIVERSITY OF WISCONSIN-MADISON

Provided to you by:

## Pear Slug (Pear Sawfly)

Phil Pellitteri, UW Insect Diagnostic Lab

Pear slugs are not true slugs (see University of Wisconsin Garden Facts XHT1040), but are larvae of the pear sawfly (*Calora cerasi*). These larvae feed and cause damage on many kinds of ornamentals and fruit trees, including cherry, cotoneaster, mountain-ash, pear, purple leaf plum and serviceberry.

**Appearance:** Pear slugs (larvae) are dark green to orange in color, tadpole-shaped and grow up to ½ inch in length. They have a shiny appearance because they cover their bodies with their own slimy, liquid waste. Pear sawflies (adults) are shiny-black, thick-bodied, stingless wasps that are ⅛ to ¼ inch long with ¼ to ⅜ inch wide wingspans.



Pear slugs are tadpole-shaped and slimy in appearance. They feed on the tissue between leaf veins giving leaves a lacy appearance.

**Damage:** Pear slugs feed on the upper surface of leaves, and typically avoid feeding on lower leaf surfaces. When feeding, pear slugs eat the tissue between leaf veins, but leave the veins themselves intact. This feeding pattern is known as skeletonization and gives leaves a lacy appearance. If feeding is extensive, leaves may brown and shrivel. In general, damage from pear slug is cosmetic and will not harm the overall health of a plant.

**Life Cycle:** Pear sawfly adults emerge from overwintering cocoons in late June through July, and subsequently insert eggs into the upper surface of leaves of preferred hosts. The eggs hatch and the resulting pear slugs (larvae) feed for two to three weeks. Larvae pupate in the soil and a second generation of adults appears and lays eggs in August through early September. The resulting larvae then feed for two to three weeks and overwinter as cocoons in the soil.

**Control:** Pear slugs often complete their feeding long before gardeners notice that leaf damage has occurred. Therefore, before considering any treatments, make sure that larvae are present on plants. Pear slugs are easily killed with insecticidal soap, neem oil, bifenthrin, carbaryl, malathion, permethrin, spinosad or other general landscape insecticide sprays. Pear slugs are not controlled by bacterial products containing *Bacillus thuringiensis* (Bt). Soil drenches also are not effective for controlling this pest.

**For more information on pear slugs (pear sawflies):** Contact your county Extension agent.

© 2010-2019 by the Board of Regents of the University of Wisconsin System doing business as the division of Cooperative Extension of the University of Wisconsin Extension. An EEO/Affirmative Action employer, University of Wisconsin Extension provides equal opportunities in employment and programming, including Title IX and ADA requirements. This document can be provided in an alternative format by calling Brian Hudelson at (608) 262-2863 (711 for Wisconsin Relay).

References to pesticide products in this publication are for your convenience and are not an endorsement or criticism of one product over similar products. You are responsible for using pesticides according to the manufacturer's current label directions. Follow directions exactly to protect the environment and people from pesticide exposure. Failure to do so violates the law.

Thanks to Diana Alfuth, Donna Henderson, Barb Larson and Scott Reuss for reviewing this document.

A complete inventory of University of Wisconsin Garden Facts is available at the University of Wisconsin-Madison Division of Extension Plant Disease Diagnostics Clinic website: <https://pddc.wisc.edu>.