



Extension

UNIVERSITY OF WISCONSIN-MADISON

Provided to you by:

Grub Control in Home Lawns

R. Chris Williamson, UW Turf and Ornamental Specialist

M. J. Ballweg, UW-Extension Sheboygan County

Listed below are trade names and common chemical names of insecticides available to homeowners for controlling white grubs in lawns.



White grub damage.

Preventative Control Products

Due to the development of new and improved insecticide chemistries, season-long residual or “preventive” controls of white grubs are now available, and are becoming the preferred management strategy. Preventative control strategies require insecticide application prior to egg lay or hatch. Timing varies depending on grub species: May/June beetle; late-May – early-June, and Japanese beetle; late-June – late-July.

Preventative Control Products				
Trade Name	Company	Common Chemical Name	Overall Toxicity	Application Timing
Season-Long Grub Control	Bayer Advanced Lawn	Imidacloprid	Low	Late-May – Late July
GrubEx	Scotts	halofenozide	Low	Late-May – Late July

Applications of preventative control products should not be made after **August 1st**!



Extension

UNIVERSITY OF WISCONSIN-MADISON

Curative Control Products

The curative control approach entails applying a control product when grubs are present, active, and causing measurable damage. Insecticides should be applied when grubs are feeding in the root zone. However, greater control can be attained if the insecticide is applied to smaller or younger grubs – usually in July for May/June beetles, or August for Japanese beetles.

Curative Control Products				
Trade Name	Company	Common Chemical Name	Overall Toxicity	Application Timing
24 Hour Grub Killer	Bayer	trichlorfon	Low	Mid-May – Early October
Sevin	Numerous	carbaryl	Low	Mid-May – Early October
Diazinon	Numerous	diazinon	High	No longer recommended.
Dursban	Numerous	chlorpyrifos	High	No longer recommended

Remember that grubs feed in the root zone. Thus the control product must reach them in order to be effective. Therefore, regardless of insecticide brand or formulation, you **must** water the insecticide into the root zone to achieve maximum effectiveness. Apply at least ½ inch of irrigation or rainfall on sandy soils, and up to ¾ inch on heavier clay soils. Apply water within 24 hours after application to get the most benefit from insecticides. For spray formulations, **wash spray off grass plants and into the soil before the spray dries**. Liquid formulations of some products have shown greater efficiency.

For more information on May/June beetles and Japanese beetles:
See UW-Extension bulletins A3275 and A3714, and UW-Extension Garden Facts X1062, or contact your county Extension agent.