



Extension

UNIVERSITY OF WISCONSIN-MADISON

Provided to you by:

University of Wisconsin Pest Alert

## Ralstonia Wilt

Brian Hudelson, UW-Madison Plant Pathology

**What is Ralstonia wilt?** Ralstonia wilt (also sometimes known as Southern wilt) is a usually lethal disease that affects over 250 plants in over 40 plant families. Susceptible greenhouse-grown ornamentals include, but are not limited to, plants in the genera *Capsicum*, *Cosmos*, *Cyclamen*, *Dahlia*, *Fuchsia*, *Gerbera*, *Hydrangea*, *Impatiens*, *Lantana*, *Nasturtium* and *Pelargonium*. Vegetables such as eggplant, pepper, potato and tomato, as well as tobacco, are also susceptible. Ralstonia wilt has recently been detected in geraniums (*Pelargonium* spp.) in Wisconsin.



Yellowing and wilting characteristic of Ralstonia wilt.

### What does Ralstonia wilt look like?

Symptoms of Ralstonia wilt in geraniums are similar to those associated with bacterial blight (caused by *Xanthomonas campestris* pv. *pelargonii*). Initially, lower leaves of infected plants yellow and wilt, then die. Yellowing and death of upper leaves follow. Symptoms may initially occur on only one side of the plant. Internally, the water-conducting tissue of the plant browns, and then the entire stem rots from the inside out. Finally, infected plants die.

### Where does Ralstonia wilt come from?

Ralstonia wilt is caused by the bacterium *Ralstonia solanacearum* (formerly *Pseudomonas solanacearum*). This bacterium is commonly found in tropical, sub-tropical and warm temperate climates, but is not believed to survive cold temperatures such as those typical of Wisconsin winters. The bacterium can be moved in symptomless plants or cuttings, or in contaminated soil and plant debris (where the pathogen can remain dormant for many years). Several subgroups (i.e., races or biovars) of *R. solanacearum* have been recognized, each with a different host range. *R. solanacearum* race 3, biovar 2 is of particular concern it causes a serious disease of potato called brown rot. In addition, this race/biovar has been listed as a select agent in the Agriculture Bioterrorism Act of 2002 and is considered to have potential to be developed as a bioterrorist weapon.



Extension

UNIVERSITY OF WISCONSIN-MADISON

**How do I save plants with Ralstonia wilt?** There are no known treatments that will save plants infected with Ralstonia wilt. If you believe your plants are suffering from this disease, immediately contact your local department of agriculture or county Extension agriculture or horticulture agent to arrange for confirmatory testing (see below for information on where to submit samples in Wisconsin). If your plants test positive for *R. solanacearum* race 3, biovar 2 the U.S. Animal and Plant Health Inspection Service (APHIS) must be notified and this organization will provide guidance on proper disposal of contaminated plants and decontamination of greenhouses.

**How do I avoid problems with Ralstonia wilt in the future?** Start by purchasing and growing pathogen-free plant cuttings. Keep plants from different suppliers physically separated by at least 4 ft. to minimize the risk of cross contamination should a shipment of plants prove to be contaminated. Because *R. solanacearum* is easily moved with soil or water, minimize splashing or any other movement of water or soil from plant to plant when watering. When taking cuttings or trimming plants, be sure to clean cutting tools between cuts by dipping them in a 10% bleach solution, ammonia or alcohol (spray disinfectants that contain at least 70% alcohol can also be used). Also wash your hands frequently when handling plants to minimize the possibility of moving the bacterium by touch. Remove and destroy weeds or weed debris as these can harbor the pathogen. Finally, do not grow plants in a greenhouse where the disease has occurred unless it has been properly decontaminated.

**For more information on Ralstonia wilt or help in diagnosing this problem:** Contact Brian Hudelson, Plant Disease Diagnostics Clinic, University of Wisconsin-Madison/Extension, 1630 Linden Drive, Madison, WI 53706-1598 [phone: (608) 262-2863, fax: (608) 263-2626, email: [bdh@plantpath.wisc.edu](mailto:bdh@plantpath.wisc.edu)].

© 1999-2019 by the Board of Regents of the University of Wisconsin System doing business as the division of Cooperative Extension of the University of Wisconsin Extension.

An EEO/Affirmative Action employer, University of Wisconsin Extension provides equal opportunities in employment and programming, including Title IX and ADA requirements. This document can be provided in an alternative format by calling Brian Hudelson at (608) 262-2863 (711 for Wisconsin Relay).

References to pesticide products in this publication are for your convenience and are not an endorsement or criticism of one product over similar products. You are responsible for using pesticides according to the manufacturer's current label directions. Follow directions exactly to protect the environment and people from pesticide exposure. Failure to do so violates the law.

Thanks to Alemu Mengistu, Julie Tans-Kersten and Laurie Weiss for reviewing this document.

A complete inventory of University of Wisconsin Garden Facts is available at the University of Wisconsin-Madison Division of Extension Plant Disease Diagnostics Clinic website: <https://pddc.wisc.edu>.